



**Contacts:**

Hillary Wasserman  
Senior Associate  
[HWasserman@GroupGordon.com](mailto:HWasserman@GroupGordon.com)  
+1 732-778-1896

Becky Bunn, MSc  
Public Relations Manager  
[Becky.Bunn@IASLC.org](mailto:Becky.Bunn@IASLC.org)  
+1 720-254-9509

## **Treatment Based On BRCA1 Expression Levels Does Not Increase Survival Rate of Stage II and III Non-Small Cell Lung Cancer N+ Resected Patients**

Yokohama, Japan - October 17, 2017 - New research shows that treating stage II and III non-small cell lung cancer (NSCLC) N+ resected patients with customized chemotherapy (CT) treatment based on their specific BRCA1 expression levels, as opposed to providing the standard treatment, did not increase overall survival rates among those patients who received individualized CT treatment. However, BRCA1 expression levels could be prognostic and treatment could achieve different outcomes. Dr. Bartomeu Massuti of Alicante University Hospital in Spain on behalf the Spanish Lung Cancer Cooperative Group (GECP-SCLG) presented his findings today at the International Association for the Study of Lung Cancer (IASLC) 18<sup>th</sup> World Conference on Lung Cancer (WCLC) in Yokohama, Japan.

For patients with resected NSCLC with lymph node involvement, postoperative platinum-based CT is the standard of care. BRCA1 is a DNA repair factor that is mainly involved in the repair of double strand DNA breaks and may also act as a differential regulator of response to cisplatin (Cis) and antimicrotubule agents. BRCA1 efficiency enhances resistance to Cis, and loss of BRCA1 function is associated with sensitivity to DNA-damaging CT. Therefore, the research team sought to determine if differential CT treatment based on knowledge of patients' BRCA1 levels could lead to higher survival rates and if cisplatin could be avoided in a pre-defined subgroup.

The researchers hypothesized that the five year survival rate of the standard care control group (45%) could increase by 20% in an intervention group that received CT compatible with either their low, medium or high BRCA1 expression levels. While those in the control group received the standard Cis-Docetaxel, those in the experimental group either received Cis-Gemcitabine (low BRCA1), Cis-Doc (intermediate BRCA1) or Docetaxel (high BRCA 1).

After median follow-up of 53 months, the OS among the control group was 69.3 months and among the experimental group 82.3 months, ranging from 74 months (low BRCA) to 80.5 months (intermediate BRCA) and 80.2 months (high BRCA).

Their findings suggest that survival rates do not increase among patients who receive chemotherapy (CT) treatment based on their BRCA1 levels. However, the researchers did find higher rates of survival than expected in patients with lymph node involvement, and also determined that in the case of high levels of BRCA1, CT treatment with Docetaxel (but without cisplatin) is not detrimental. Also for patients with low BRCA1 levels, Cisplatin-Gemcitabine could be superior to Cisplatin-Docetaxel.

“The findings from our study do not allow changing the current standard of treatment, but this trial validates BRCA1 as a prognostic factor and selection of chemotherapy treatment based on BRCA1 expression levels could be an option to improve outcomes, avoiding in some cases the use of cisplatin and its toxicity,” said Dr. Massuti. “Having this evidence to back up our treatment decisions is essential to providing the best care possible and also opens new opportunities for research in this setting.”

### **About the WCLC**

The World Conference on Lung Cancer (WCLC) is the world’s largest meeting dedicated to lung cancer and other thoracic malignancies, attracting over 6,000 researchers, physicians and specialists from more than 100 countries. The goal is to disseminate the latest scientific achievements; increase awareness, collaboration and understanding of lung cancer; and to help participants implement the latest developments across the globe. Organized under the theme of “Synergy to Conquer Lung Cancer,” the conference will cover a wide range of disciplines and unveil several research studies and clinical trial results. For more information, visit <http://wclc2017.iaslc.org/>.

### **About the IASLC**

The International Association for the Study of Lung Cancer (IASLC) is the only global organization dedicated to the study of lung cancer and other thoracic malignancies. Founded in 1974, the association's membership includes more than 6,500 lung cancer specialists across all disciplines in over 100 countries, forming a global network working together to conquer lung and thoracic cancers worldwide. The association also publishes the Journal of Thoracic Oncology, the primary educational and informational publication for topics relevant to the prevention, detection, diagnosis and treatment of all thoracic malignancies. Visit [www.iaslc.org](http://www.iaslc.org) for more information.

# # #